

Climate Impacts of Methane Losses from Modern Natural Gas & Petroleum Systems


The climate implications of increased natural gas production and use, particularly substitution of coal by natural gas in the power sector has been hotly debated since 2011. Because natural gas combustion is associated with half the carbon dioxide (CO₂) emissions of coal in the electric power sector, it is often assumed that a switch to natural gas power will result in lowered greenhouse gas emissions. However, this assumption does not account for methane emissions associated with producing natural gas and bringing that natural gas to market.

The past several years have seen major changes both in our understanding of the importance of reducing methane emissions as a climate mitigation measure, and the significance of natural gas and petroleum systems as a source of atmospheric methane. Here, we review the current literature to provide the most up-to-date calculations for the climate impact and magnitude of methane emissions from modern natural gas and oil development.


3.8% Methane losses from natural gas drilling to distribution based on atmospheric measurements[†]
Gross U.S. production

At this rate of loss, fuel conversion from coal to natural gas would not produce a climate benefit for another 40 years.

[†] Weighted average of methane losses from natural gas and petroleum systems relative to gross natural gas production from natural gas and oil wells (see page 2). Assumes default gas composition 88% methane. *Adapted from Alvarez et al. (2012)² using AR5 radiative efficiency for methane³ and 3.8% loss rate.



Rapid methane mitigation combined with CO₂ mitigation is society's best chance for stabilizing the climate. Climate scientists estimate a threshold of 1.8° - 2° C over pre-industrial levels. Additional warming beyond this threshold is associated with the rapid onset of a climate never before experienced by humans which we may not be able to adapt to fast enough.³



Natural Gas is not a Bridge Fuel to a Lower Carbon Future. Updated Technology Warming Potentials (TWP)² using the most recent IPCC findings³ indicate an emission limit of 2.8% of natural gas production for conversion from coal. **Loss rates above 2.8% negate any near-term climate benefits associated with lower carbon dioxide emissions during fuel combustion.**

- Methane emissions inferred from atmospheric concentrations reflect the weighted average emissions attributed to onshore natural gas and petroleum systems as reported in 13 atmospheric studies (see emissions discussion, page 2);
- Methane emissions inferred from atmospheric sampling exceed the emissions limit for substitution of coal by at least 40%;
- Conversion to natural gas in vehicles to replace gasoline and diesel have much lower TWPs and already exceed emission limits based on EPA emission estimates.



Methane Emissions from Petroleum and Natural Gas Systems.

- Methane, the primary component of natural gas, is produced from both natural gas (non-associated) and oil (associated) wells;
- Fugitive emissions (unintentional) and vented (intentional) emissions occur throughout both supply chains;
- U.S. EPA estimates methane emissions from oil and gas activity to be 7,589 Gg methane (10^9 g) per year, or roughly 27% of all 2013 methane emissions in the U.S.¹, **BUT...**

Upstream: Exploration, Drilling, and Production

Midstream: Natural gas Gathering & Processing; Petroleum Transport

Downstream: Natural gas Transmission/Storage & Distribution; Petroleum Refining



Atmospheric methane measurements indicate that actual emissions from these systems are at least a factor of 2 - 2.5 higher than what is reported in EPA inventory.

Our comparison of EPA inventory estimates and emissions inferred from atmospheric sampling focus solely on field measurement studies published since 2011⁴⁻¹⁶ and emissions attributable specifically to natural gas and petroleum sectors.

Based on these measurements, weighted by the study region's percent of gross domestic production, **we estimate that real-world 2013 U.S. methane emissions from petroleum and natural gas systems were 16,141 to 18,952 Gg/y**, compared to EPA's estimate of 7,589 Gg/y.

13 Atmospheric measurement studies published between 2011 and 2015

References

1. USEPA (2015). Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2013. EPA 430-R-13-001. US Environmental Protection Agency Washington DC.
2. Alvarez, R. A., et al. (2012). Greater focus needed on methane leakage from natural gas infrastructure. *PNAS*. dx.doi.org/10.1073/pnas.1202407109
3. IPCC (2013). Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
4. Petró, G., et al. (2012). Hydrocarbon emissions characterization in the Colorado Front Range: A pilot study. *J Geophys Res*. dx.doi.org/10.1029/2011JD016360
5. Wennberg et al. (2012). On sources of methane to the Los Angeles atmosphere. *ES&T*. dx.doi.org/10.1021/es301138y
6. Karion, A., et al. (2013). Methane emissions estimate from airborne measurements over a western United States natural gas field. *Geophys Res Lett*. dx.doi.org/10.1002/grl.50811
7. Miller, SM, et al. (2013). Anthropogenic emissions of methane in the United States. *PNAS*. dx.doi.org/10.1073/pnas.1314392110
8. Caulton D, et al. (2014). Toward a Better Understanding and Quantification of Methane Emissions from Shale Gas Development. *PNAS*. dx.doi.org/10.1073/pnas.1316546111
9. Petron, G., et al. (2014). A new look at methane and non-methane hydrocarbon emissions from oil and natural gas operations in the Colorado Denver-Julesburg basin. *J Geophys Res*. dx.doi.org/10.1002/2013JD021272
10. Schneising, O, et al. (2014) Remote sensing of fugitive methane emissions from oil and gas production in North American tight geologic formations. *Earth's Future*. dx.doi.org/10.1002/2014EF000265
11. Lavoie et al. (2015). Aircraft-based measurements of point source methane emissions in the Barnett Shale Basin. *ES&T*. dx.doi.org/10.1021/acs.est.5b00410
12. Lyon et al. (2015). Constructing a spatially resolved methane emission inventory for the Barnett Shale region. *ES&T*. dx.doi.org/10.1021/es506359c
13. Marchese et al. (2015). Methane emissions from United States natural gas gathering and processing. *ES&T*. dx.doi.org/10.1021/acs.est.5b02275
14. McKain et al. (2015). Methane emissions from natural gas infrastructure and use in the urban region of Boston, Massachusetts. *PNAS*. dx.doi.org/10.1073/pnas.1416261112
15. Peisch, J, et al. (2015). Quantifying sources of methane using light alkanes in the Los Angeles basin, California. *Journal of Geophysical Research: Atmospheres* **118**: 4974-4990
16. Zimmerle et al. (2015). Methane emissions from the natural gas transmission and storage system in the United States. *ES&T*. dx.doi.org/10.1021/acs.est.5b01669